| Sturgeon Reservoir | Bank Anglers | | Subs | Leg Rel | Leg Kept | Over- | 2001 Cum Leg Kept | Boat Anglers | | Subs | Leg Rel | _ | Over- 2001 Cu size Leg Kep | |
|------------------------------------|------------------------|--------------|-------------|-------------|-------------|-------|----------------------|-----------------|--------------|-------------|-------------|------------|-------------------------------|----|
| Bonneville Pool | 57 | | 21 | 1 | 2 | | 2 | 20 | | 31 | | · | 3 | 5 |
| The Dalles Pool | | | 1 | 1 1 | | 1 | 22 13 | | | 3 | 6 | 7 | | |
| John Day Pool | 5 | | | | | | • | 4 | | - 4.4 | | | | 40 |
| Total | 97 | | 22 | 1 | 3 | | 3 | 46 | | 44 | | 3 | 9 | 12 |
| Salmonids | Bank Anglers | Wild Kept | Wild Rel | Hat Kept | Hat Rel | | 2001 Cum Leg Kept | Boat Anglers | Wild Kept | Wild Rel | Hat Kept | Hat Rel | 2001 Cu Leg Kep | |
| Co | inook ho eelhead | | | | | | | | | | | | | |
| The Dalles Pool Ch Co | inook | | | 1 | | | 3 | 28 | | 2 | 11 | 1 | 15 | 18 |
| Ste | elhead | | | 1 | | | 3 | | | 2 | 11 | 1 | 15 | 18 |
| John Day Pool Ch Co | 2 inook ho | | | | | | | 8 | | 4 | 3 | | 3 | 3 |
| Ste | elhead | | | | | | | | | 4 | 3 | | 3 | 3 |
| Walleye | Bank Anglers | | Leg Rel | Leg Kept | | | 2001 Cum Leg Kept | Boat Anglers | | Leg Rel | Leg Kept | | 2001 Cu Leg Kep | |
| Bonneville Pool | | | | | | | | | | | | | | |
| The Dalles Pool John Day Pool | | | | | | | | 5 4 | | | 1 | | 1 | 1 |
| Bass | Bank Anglers | | Rel | Kept | | | 2001 Cum Leg Kept | Boat Anglers | | Rel | Kept | | 2001 Cu Leg Kep | |
| Bonneville Pool The Dalles Pool | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | |

¹ Quantities include both Washington and Oregon samples.

John Day Pool

Bonneville: Sturgeon effort was 108% of the 2-week average and CPUE was 0.71 fish per angler. There was no in-sample effort for other species.

The Dalles: Sturgeon effort was 90% of the 2-week average and CPUE was 0.32 fish per angler. Salmonid effort was 125% and CPUE was 0.45. Walleye effort was 100% and CPUE was 0.20. There was no in-sample bass effort.

John Day: Sturgeon effort was 78% of the 2-week average with no reported catch. Salmonid effort was 200% and CPUE was 0.70. Walleye effort was 200% with no reported catch. There was no in-sample effort for bass.

WDFW salmonid sport sampling summary for Washington Columbia River tributaries - Jan. 8-14.

| | Bank | Bank Chinook | | Coho | | Steelhead | | | Boat | Chinook | | Coho | | Steelhead | | | |
|---------------|---------|--------------|-------|--------|-------|-----------|------|-------|---------|---------|-------|--------|-------|-----------|------|-------|--|
| Area | Anglers | Adults | Jacks | Adults | Jacks | Kept | Rel. | Other | Anglers | Adults | Jacks | Adults | Jacks | Kept | Rel. | Other | |
| Cowlitz River | 9 | 0 | 0 | 0 | 0 | 5 | 0 | 0 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Lewis River | 22 | CLOSED | | 0 | 0 | 0 | 0 | 0 | 10 | CLOSED | | 0 | 0 | 0 | 0 | 0 | |